

Measures

- #47 Care Plan
- #134 Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan
- #160 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis
- #205 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis
- #226 Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention
- #338 HIV Viral Load Suppression
- #339 Prescription of HIV Antiretroviral Therapy
- #340 HIV Medical Visit Frequency

Reporting Instructions

- (1) Identify patients with the following criteria:
 - Patient sample criteria for the HIV/AIDS Measures Group are patients aged 13 years and older with a specific diagnosis of HIV/AIDS accompanied by a specific patient encounter
One of the following diagnosis codes indicating HIV/AIDS:
B20, Z21
Accompanied by:
One of the following patient encounter codes: 99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, G0402
- (2) Report all applicable measures within the measures group for at least 20 patients.
 - The majority of the patients reported must be Medicare Part B Fee-for-service (FFS) patients, including Medicare Secondary and Railroad Medicare.
 - For each measure, the measure's clinical performance must be satisfied for at least 1 patient. [Note: An asterisk (*) indicates that the measure's clinical performance is satisfied.]

For additional information or clinical rationale for measures, review the CMS Measure Specification for 2016 PQRS. All measures group Measure Specifications can be found on the NJII Member Portal > Registry > Getting Started section.

2016 PQRS HIV/AIDS Measures Group

Patient Information

First Name

Middle Initial

Last Name

Date of Birth

Visit Date

Medical Record Number

Gender

Male

Female

Medicare FFS Patient

Yes

No

Select Measures Performance

(page 1 of 3)

* Recommended clinical performance satisfied

^ Patient is excluded from measure's performance

Select one (1) clinical action for each measure below, where applicable:

#47: Care Plan

Patients (65 years or older) who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but patient did not wish or was unable to name a surrogate decision maker or provide an advance care plan.

Advance Care Plan or surrogate decision maker documented in the medical record *

Advance Care Planning discussed and documented, patient did not wish to name surrogate decision maker *

Advance care planning not documented, (No reason documented)

Select Measures Performance

(page 2 of 3)

#134 Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan

Patients screened for clinical depression on the date of the encounter using an age appropriate standardized tool AND, if positive, a follow-up plan is documented on the date of the positive screen

Screening for clinical depression is documented as being positive AND a follow-up plan is documented*

Screening for clinical depression documented as negative (follow-up plan not required)*

Clinical depression screening not documented (Patient not eligible)^

Screening for clinical depression documented as positive, no follow-up plan documented (Reason Documented)^

Clinical depression screening not documented (No Reason Documented)

Screening for clinical depression documented as positive, no follow-up plan documented (No Reason Documented)

#160 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis

Patients with a diagnosis of dementia who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan

Prescribed*

Not Prescribed (Medical reason documented)^

Not Prescribed (CDC4+ cell count \geq 200)^

Not Prescribed (No reason documented)

#205 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis

Patients with chlamydia, gonorrhea, and syphilis screenings performed at least once since the diagnosis of HIV infection

Results documented for all 3 screenings*

Screening results not documented (Patient refusal)^

Screening results not documented (No reason documented)

Need more information? Connect with NJII PQRS Registry
973-642-4055 • pqrs@njii.com • www.njii.com/pqrs

Select Measures Performance

(page 3 of 3)

#226 Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention

Patients screened for tobacco use at least once within 24 months AND received tobacco cessation counseling intervention if identified as a tobacco user

Identified as tobacco user and cessation intervention performed*

Screened for tobacco use and identified as non-user of tobacco*

Not screened for tobacco use (Medical reason documented)^

Identified as tobacco user and cessation intervention NOT performed

Not screened for tobacco use (No reason documented)

#338 HIV Viral Load Suppression

Number of patients with a HIV viral load less than 200 copies/mL at last viral load test

Viral load < 200 copies/mL*

Viral load \geq 200 copies/mL, or viral load not performed

#339 Prescription of HIV Antiretroviral Therapy

Number of patients prescribed HIV antiretroviral therapy during the reporting period

Antiretroviral therapy prescribed*

Antiretroviral therapy not prescribed

#340 HIV Medical Visit Frequency

Number of patients who had at least one medical visit in each 6 month period of the 24 month measurement period, with a minimum of 60 days between medical visits

Patient had at least one medical visit in each 6 month period of the 24 month measurement period, with a minimum of 60 days between medical visits*

Patient did not have at least one medical visit in each 6 month period of the 24 month measurement period, with a minimum of 60 days between medical visits