

2016 PQRS

Recommended Measures for: Colorectal Surgery

Measures Groups

Reporting Requirements:

- Choose one (1) Measures Group
- Report on a minimum of 20 eligible patients (at least 11 must be Medicare Part B FFS patients)
- The numerator requirement must be met at least once for each reported measure

General Surgery

- #130: Documentation of Current Medications in the Medical Record
- #226: Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention
- #354: Anastomotic Leak Intervention
- #355: Unplanned Reoperation within the 30 Day Postoperative Period
- #356: Unplanned Hospital Readmission within 30 Days of Principal Procedure
- #357: Surgical Site Infection (SSI)
- #358: Patient-Centered Surgical Risk Assessment and Communication

Preventive Care Measures Group

- #39: Screening for Osteoporosis for Women Aged 65 - 85 Years of Age
- #48: Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older
- #110: Preventive Care and Screening: Influenza Immunization
- #111: Pneumonia Vaccination Status for Older Adults
- #112: Breast Cancer Screening
- #113: Colorectal Cancer Screening
- #128: Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan
- #134: Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan
- #226: Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention
- #431: Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling

Individual Measures

Reporting Requirements:

- Report 9 measures across 3 National Quality Strategy (NQS) domains
- For each measure, report the quality action performed for at least 50% of the Medicare Part B FFS patients who are eligible for the measure.
- Include at least 1 cross-cutting measure **CC**
- The numerator requirement must be met at least once for each reported measure.

2016 PQRS

Recommended Measures for: Colorectal Surgery



Measures by Quality Domain

Communication and Care Coordination

#47: Care Plan **CC**

#185: Colonoscopy Interval for Patients with a History of Adenomatous Polyps – Avoidance of Inappropriate Use

#320: Appropriate Follow-Up Interval for Normal Colonoscopy in Average Risk Patients

Community/Population Health

#110: Preventive Care and Screening: Influenza Immunization **CC**

#111: Pneumonia Vaccination Status for Older Adults **CC**

#134: Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan **CC**

#226: Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention **CC**

#317: Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented **CC**

Effective Clinical Care

#425: Photodocumentation of Cecal Intubation

Patient Safety

#21: Perioperative Care: Selection of Prophylactic Antibiotic – First OR Second Generation

#22: Perioperative Care: Discontinuation of Prophylactic Parenteral Antibiotics (Non-Cardiac Procedures)

#23: Perioperative Care: Venous Thromboembolism (VTE) Prophylaxis (When Indicated in ALL Patients)

#130: Documentation of Current Medications in the Medical Record **CC**

Person and Caregiver-Centered Experience and Outcomes

#342: Pain Brought Under Control Within 48 Hours

#358: Patient-Centered Surgical Risk Assessment and Communication

Need more information? Connect with NJII PQRS Registry
973-642-4055 • pqrs@njii.com • www.njii.com/pqrs

2016 PQRS

Recommended Measures for: Colorectal Surgery



Measures Applicability Validation (MAV) Clusters

View the [2016 PQRS MAV Process for Registry-Based Reporting of Individual Measures](#) for more details.

Cluster 20: Perioperative Care

- #21: Perioperative Care: Selection of Prophylactic Antibiotic – First OR Second Generation Cephalosporin
- #22: Perioperative Care: Discontinuation of Prophylactic Parenteral Antibiotics (Non-Cardiac Procedures)
- #23: Perioperative Care: Venous Thromboembolism (VTE) Prophylaxis (When Indicated in ALL Patients)

Cluster 22: Endoscopy and Polyp Surveillance

- #185: Colonoscopy Interval for Patients with a History of Adenomatous Polyps – Avoidance of Inappropriate Use
- #320: Appropriate Follow-Up Interval for Normal Colonoscopy in Average Risk Patients

Cluster 35: Colonoscopy Care

- #130: Documentation of Current Medications in the Medical Record **CC**
- #425: Photodocumentation of Cecal Intubation

Cluster 40: Palliative Care

- #47: Care Plan **CC**
- #134: Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan **CC**
- #342: Pain Brought Under Control Within 48 Hours

Cluster 41: General Care

- #130: Documentation of Current Medications in the Medical Record **CC**
- #134: Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan **CC**
- #226: Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention **CC**

Cluster 43: Immunization Care

- #110: Preventive Care and Screening: Influenza Immunization **CC**
- #111: Pneumonia Vaccination Status for Older Adults **CC**

Need more information? Connect with NJII PQRS Registry
973-642-4055 • pqrs@njii.com • www.njii.com/pqrs